arXiv:1103.4292v2 [math.AG] 15 Mar 2012

A UNIFORM BOUND ON THE CANONICAL DEGREE OF ALBANESE DEFECTIVE CURVES ON SURFACES

MARGARIDA MENDES LOPES AND RITA PARDINI

Abstract. Let S be a minimal complex surface of general type with irregularity $q \geq 2$ and let $C \subset S$ be an irreducible curve of geometric genus g. Assume that C is *Albanese defective*, i.e., that the image of C via the Albanese map does not generate the Albanese variety $\mathrm{Alb}(S)$; we obtain a linear upper bound in terms of K_S^2 and g for the canonical degree K_SC of C. As a corollary, we obtain a bound for the canonical degree of curves with $g \leq q-1$, thereby generalizing and sharpening the main result of [Lu].

2000 Mathematics Subject Classification: Primary 14J29; Secondary 14C20, 32C25.
Keywords: curves on irregular surfaces, uniform bound on canonical degree, irregular surface.

1. Introduction

Let S be an irregular minimal surface of general type with irregularity $q \geq 2$ and let $a\colon S \to \mathrm{Alb}(S)$ be the Albanese map. Let $C \subset S$ be a curve; we define the *Albanese defect* $\delta_{\mathrm{alb}}(C)$ to be the codimension in $\mathrm{Alb}(S)$ of $< a(C) >$, the smallest abelian subvariety of $\mathrm{Alb}(S)$ containing $a(C)$. If $\delta_{\mathrm{alb}}(C) > 0$, then we say that C is *Albanese defective.*

In this note we give a uniform linear bound for the canonical degree K_SC of an Albanese defective curve C in terms of K_S^2 and of the geometric genus g of C. For a general discussion on bounds of the canonical degree of curves see [Lu], [LM] and [Mi].

We prove the following theorem:

Theorem 1.1. *Let S be a minimal surface of general type with irregularity $q \geq 2$ and let C be an irreducible curve of S with geometric genus g such that $\delta_{\mathrm{alb}}(C) > 0$. Then:*

- *(i) if $\delta_{\mathrm{alb}}(C) \geq 2$, then $K_SC \leq K_S^2 - \delta_{\mathrm{alb}}(C)$;*
- *(ii) if $\delta_{\mathrm{alb}}(C) = 1$, then $K_SC \leq 4K_S^2 + 2g - 2$ and there is a fibration $f\colon S \to B$, such that $g(B) = 1$ and C is a component of a fibre.*

The first author is a member of the Center for Mathematical Analysis, Geometry and Dynamical Systems (IST/UTL); the second author is a member of G.N.S.A.G.A. of In.D.A.M.. This research was partially supported by FCT (Portugal) through program POCTI/FEDER and Project PTDC/MAT/099275/2008 and by italian MIUR through PRIN 2008 project "Geometria delle varietà algebriche e dei loro spazi di moduli".

If C is an Albanese defective curve, then the map $S \to A/{<}a(C){>}$ is a non constant morphism that contracts C to a point, hence one has $C^2 \leq 0$. In [LM] it is proven that if S is a (not necessarily irregular) surface of general type and $C \subset S$ is an irreducible curve of genus g with $C^2 \leq 0$ then $K_SC \leq 6c_2(S)-2K_S^2+6g-6$. So Theorem 1.1 can be viewed as a refinement of this inequality in the special case of an Albanese defective curve. Note also that the bounds in Theorem 1.1 imply $p_a(C) - g(C) \leq 2K_S^2$, namely for an Albanese defective curve the difference between the arithmetic genus and the geometric genus is bounded by a fixed quantity depending only on the surface S. Explicit examples of Albanese defective curves with several singularities are not easy to produce; a construction can be found in Example 4.4

If a curve C has geometric genus $g < q$, then $\delta_{\mathrm{alb}}(C) \geq q - g > 0$. So we have the following immediate consequences of Theorem 1.1:

Theorem 1.2. *Let S be a minimal surface of general type with irregularity $q \geq 2$ and let C be an irreducible curve of S with geometric genus $g < q$. Then:*

- *(i) if $g \leq q - 2$, then $K_SC \leq K_S^2 - (q - g)$;*
- *(ii) if $g = q - 1$, then either $K_SC \leq K_S^2 - 2$, or there is a fibration $f \colon S \to B$, where $g(B) = 1$, and C is a component of a fibre. In this case $K_SC \leq 4K_S^2 + 2q - 4$.*

Corollary 1.3. *Let S be a surface of general type with irregularity $q \geq 2$. Then:*

- *(i) the cohomology classes in $H^2(S, \mathbb{Z})$ of the irreducible curves C of geometric genus $< q$ are a finite set;*
- *(ii) if S has no nontrivial morphism onto a curve of genus > 0, then S contains finitely many curves of geometric genus $< q$.*

Theorem 1.2 and Corollary 1.3, (ii) generalize a result of S.Y. Lu ([Lu]) for curves of geometric genus ≤ 1.

The first part of the proof of Theorem 1.1 uses the same ideas as the first part of the proof in [Lu]. However the remainder of the proof is different and shorter, and it does not require a deep analysis of the behaviour of ramification of the Albanese map. This simplification is made possible by the use of two extra ingredients: the smoothness of bicanonical curves (see, e.g., [Ci]) and the Severi inequality for minimal irregular surfaces of maximal Albanese dimension ([Pa]).

If F is a smooth fiber of a fibration $S \to E$, with E an elliptic curve, then $K_SF = 2g(F) - 2$; since there are surfaces with infinitely many fibrations onto an elliptic curve with $g(F) \to \infty$ (see [MP1], Example 2.1.2), any bound for the canonical degree of Albanese defective curves must be $\geq 2g - 2$, and so the term $2g - 2$ in Theorem 1.1 (ii) seems to be optimal. However, the bound is possibly not sharp; if it were, then several inequalities used in the proof should be equalities, all at the same time. In particular the Severi inequality should be an equality. Conjecturally this happens only for $q = 2$, but it has been proven only under the assumption that the canonical class of the surface be ample ([Ma], cf. also [MP2]); in the general case it has been an open question for many years. Also, the coefficient 4 in the term in function of K_S^2 seems too large; still, we find it intriguing that the different proof for the case of elliptic curves in [Lu] also gives the same coefficient 4.

In §4 we compute K_SC for some examples of Albanese defective curves C.

Notation and conventions: All varieties are complex projective; a *surface* is a smooth projective surface. The standard notation for the invariants of a surface S is used: $q(S) := h^0(\Omega^1_S) = h^1(\mathcal{O}_S)$ is the *irregularity*, $p_g(S) := h^0(K_S)$ is the *geometric genus* and $\chi(\mathcal{O}_S) = 1+p_g(S)-q(S)$ is the (holomorphic) Euler-Poincaré characteristic and $c_1(S)$, $c_2(S)$ are the Chern classes of the tangent bundle of S. $\chi_{\text{top}}(X)$ denotes the topological Euler characteristic of a space X.

Recall that $K_S^2 = c_1(S)^2$, $c_2(S) = \chi_{\text{top}}(S)$ and that the Noether formula $K_S^2 + c_2(S) = 12\chi(\mathcal{O}_S)$ holds.

An effective divisor D on S is *k-connected* if for every decomposition $D = A+B$, with $A, B > 0$ one has $AB \geq k$. We sometimes refer to effective divisors as "curves". As usual we denote by $p_a(D)$ the arithmetic genus of a curve and by $g(D)$ the geometric genus of an irreducible curve.

The Jacobian of a smooth curve B is denoted by $J(B)$.

2. Preliminaries

In this section we collect several facts that will be of use in the proof of Theorem 1.1.

We recall the following corollary of Arakelov's theorem.

Proposition 2.1 (see [Be2], Corollaire)**.** *Let $f\colon S \to B$ be a fibration of a smooth minimal surface S onto a smooth curve B of genus b, with general fibre of genus $h \geq 2$. Then*

$$8(h-1)(b-1) \leq K_S^2.$$

Next we give a lower bound for the canonical degree of some curves of S.

Lemma 2.2. *Let S be a smooth minimal surface of general type and let D be a curve such that $K_S - D > 0$. Then*

$$K_SD \geq q - \delta_{\text{alb}}(D).$$

Proof. By the 2-connectedness of canonical divisors we have $D(K_S-D) \geq 2$, namely $K_SD \geq D^2 + 2$. If D is irreducible, by the adjunction formula we have:

$$2K_SD \geq 2p_a(D) \geq 2g(D) \geq 2(q - \delta_{\text{alb}}(D)).$$

In general, the inequality follows by observing that if $D_1, \dots D_r$ are the distinct irreducile components of D, then $K_SD \geq \sum_i K_SD_i$ and $q - \delta_{\text{alb}}(D) \leq \sum_i(q - \delta_{\text{alb}}(D_i))$. $\square$

We will need also the following:

Proposition 2.3. *Let S be a surface and let $f\colon S \to B$ be a relatively minimal fibration with general fibre F of genus $g \geq 1$. Let F_0 be a singular fibre of f and C a component of F_0 appearing with multiplicity one. If N_C is the normalization of C then*

$$\chi_{\mathrm{top}}(F_0) - \chi_{\mathrm{top}}(F) \geq \frac{1}{2}K_S C + \chi(\mathcal{O}_{N_C}).$$

Furthermore if $F_0 - C > 0$ then the inequality is strict.

Proof. Write $F_0 = C + G + R$, where $C + G$ is the support of F_0 and R is an effective (possibly 0) divisor with support contained in G.

First we look at the term $-\chi_{\mathrm{top}}(F)$. Recall that

$$-\chi_{\mathrm{top}}(F) = -2\chi(\mathcal{O}_F) = K_S F + F^2,$$

where the last equality is the adjunction formula.

On the other hand, since $F_0 = C + G + R$,

$$K_S F + F^2 = K_S(C+G) + (C+G)^2 + 2(C+G)R + K_S R + R^2.$$

By the numerical properties of fibres (see, e.g., [Be1]), $RF = 0$, i.e., $R^2 + R(C+G) = 0$. Since we are assuming that $f\colon S \to B$ is a relatively minimal fibration, $K_S R \geq 0$ and so we obtain

$$-\chi_{\mathrm{top}}(F) = K_S F + F^2 \geq K_S C + C^2 + K_S G + G^2 + 2CG + (C+G)R.$$

Now we turn to $\chi_{\mathrm{top}}(F_0) = \chi_{\mathrm{top}}(C+G)$. By the additivity of the topological characteristic, one has $\chi_{\mathrm{top}}(C+G) = \chi_{\mathrm{top}}(C) + \chi_{\mathrm{top}}(G) - d$, where d is the number of intersection points of C and G. Note that $d \leq CG$.

By the proof of Lemme VI.5 of [Be1] given a reduced curve D, one has

$$\chi_{\mathrm{top}}(D) \geq \chi(\mathcal{O}_D) + \chi(\mathcal{O}_{N_D}),$$

where N_D is the normalization of D. In particular, for any reduced curve D one has $\chi_{\mathrm{top}}(D) \geq 2\chi(\mathcal{O}_D)$.

So $\chi_{\mathrm{top}}(C) + \chi_{\mathrm{top}}(G) - d \geq \chi(\mathcal{O}_C) + \chi(\mathcal{O}_{N_C}) + 2\chi(\mathcal{O}_G) - d$, where N_C is the normalization of C.

This can be written, by the adjunction formula,

$$\chi_{\mathrm{top}}(C) + \chi_{\mathrm{top}}(G) - d \geq -\frac{1}{2}(K_S C + C^2) + \chi(\mathcal{O}_{N_C}) - K_S G - G^2 - d.$$

Then $\chi_{\mathrm{top}}(F_0) - \chi_{\mathrm{top}}(F) \geq -\frac{1}{2}(K_S C + C^2) + \chi(\mathcal{O}_{N_C}) - K_S G - G^2 - d + K_S C + C^2 + K_S G + G^2 + 2CG + (C+G)R =$
$= \frac{1}{2}(K_S C + C^2) + \chi(\mathcal{O}_{N_C}) + 2CG + (C+G)R - d.$

Now $\frac{1}{2}(K_S C + C^2) + \chi(\mathcal{O}_{N_C}) + 2CG + (C+G)R - d =$

$= \frac{1}{2}K_SC + \chi(\mathcal{O}_{N_C}) + \frac{1}{2}(C^2 + CG + CR) + (CG - d) + \frac{1}{2}G(C + R) + \frac{1}{2}R(C + G)$.

From $CF = 0$ we obtain $C^2 + CG + CR = 0$. On the other hand the fibre F_0 is not a multiple fibre, because C appears with multiplicity 1 on F_0. Thus F_0 is 1-connected and so, if $G + R \neq 0$, then $\frac{1}{2}(G(C + R) + R(C + G)) > 0$. Since also $CG - d \geq 0$ we obtain

$$\chi_{\mathrm{top}}(F_0) - \chi_{\mathrm{top}}(F) \geq \frac{1}{2}K_SC + \chi(\mathcal{O}_{N_C})$$

and that if $C \neq F_0$, then the inequality is strict.

□

3. The proof of Theorem 1.1

We denote by $a\colon S \to \mathrm{Alb}(S)$ the Albanese map, we set

$$A_C := \mathrm{Alb}(S)/{<}a(C){>}$$

and we denote by $a_C\colon S \to A_C$ the induced map. Observe that A_C is an abelian variety of dimension $\delta_{\mathrm{alb}}(C)$ and that a_C is a non constant morphism that contracts C to a point.

Step 1. *If $a_C(S)$ is a surface, then $K_SC \leq K_S^2 - \delta_{\mathrm{alb}}(C)$.*
Let α, β be the pull back on S of two general 1-forms of A_C and let G be the divisor of zeros of $\alpha \wedge \beta$. Then $G = C + R$ with R is an effective divisor whose image via a_C spans A_C, hence $\delta_{\mathrm{alb}}(R) \leq q - \delta_{\mathrm{alb}}(C)$. So we have

$$K_S^2 = K_SG = K_SC + K_SR \geq K_SC + \delta_{\mathrm{alb}}(C),$$

where the last inequality follows by Lemma 2.2.

So from now we may assume that $a_C(S)$ is a curve B.

Step 2. *B is smooth of genus $\delta_{\mathrm{alb}}(C)$ and a_C has connected fibers.*
The map a_C factors through an irrational pencil $f_C\colon S \to \Gamma$, where Γ is a smooth curve of genus $\gamma \geq \delta_{\mathrm{alb}}(C)$. On the other hand, $H^0(\omega_\Gamma)$ pulls back to a γ-dimensional subspace of $H^0(\Omega_S^1)$ whose elements restrict to 0 on C, so $\gamma \leq \delta_{\mathrm{alb}}(C)$; therefore $\gamma = \delta_{\mathrm{alb}}(C)$ and the natural map $J(\Gamma) \to A_C$ is an isogeny. By the universal property of the Albanese variety, the map $\mathrm{Alb}(S) \to J(\Gamma)$ induced by f_C descends to a map $A_C \to J(\Gamma)$ such that $A_C \to J(\Gamma) \to A_C$ is the identity, hence $J(\Gamma) \to A_C$ is an isomorphism and B is smooth of genus $\delta_{\mathrm{alb}}(C)$.

Step 3. *If $\delta_{\mathrm{alb}}(C) > 1$, then $4(\delta_{\mathrm{alb}}(C) - 1)K_SC \leq K_S^2$.*
We denote by F a general fiber of f_C, which is a smooth curve of genus $g(F) \geq 2$ since S is of general type. By the adjunction formula $K_SF =$

$2g(F)-2$. Since K_S is nef, $K_SC \le K_SF$ and so, by Proposition 2.1 and Step 2, we obtain

$$4(\delta_{\mathrm{alb}}(C)-1)K_SC \le 8(g(F)-1)(\delta_{\mathrm{alb}}(C)-1) \le K_S^2.$$

Step 4. *If* $\delta_{\mathrm{alb}}(C)=1$, *then* $K_SC \le 4K_S^2+2g-2$.
In this case A_C is an elliptic curve and by Step 2 the map a_C has connected fibers; denote by F_0 the fiber of a_C which contains C.

If F_0 is non singular then $K_SF_0 = K_SC = 2g-2$ and the inequality is trivially satisfied.

So assume F_0 is singular.

Suppose that C appears in F_0 with multiplicity ≥ 2. Since $|2K_S|$ is base point free (see, e.g., [Ci]), we can consider a smooth curve D in $|2K_S|$ meeting C transversally. Let $m := K_SF_0$ and let $D \to A_C$ be the cover of degree $2m$ induced by a_C. Then one has $6K_S^2 = 2g(D)-2$ by the adjunction formula and, by the Hurwitz formula, $2g(D)-2 = \deg R$, where R is the ramification divisor of $D \to A_C$. Since C appears with multiplicity ≥ 2 in F_0, we obtain $\deg R \ge 2K_SC$. So $6K_S^2 \ge 2K_SC$, i.e. $3K_S^2 \ge K_SC$.

Suppose now that C appears in F_0 with multiplicity 1 and let F denote a general fibre of f. By Proposition 2.3,

$$\chi_{\mathrm{top}}(F_0) - \chi_{\mathrm{top}}(F) \ge \frac{1}{2}(K_SC)+1-g.$$

Then, by the formulas for c_2 of a fibered surface (see, e.g., Lemme VI.4 of [Be1]), we have $c_2(S) \ge \chi_{\mathrm{top}}(F_0) - \chi_{\mathrm{top}}(F)$.

Notice that the Albanese image of S is a surface, since $q \ge 2$ by assumption and on a surface with Albanese dimension 1 the Albanese defect of any curve is either equal to 0 or to q. Hence by the Severi inequality ([Pa]) we have $c_2(S) \le 2K_S^2$. So one obtains $\frac{1}{2}(K_SC)+1-g \le 2K_S^2$, i.e. $K_SC \le 4K_S^2+2g-2$.

Step 5. *end of proof.*
We only have to show that in the situation of Step 3 one has $K_SC \le K_S^2 - \delta_{\mathrm{alb}}(C)$.

Assume for contradiction that $K_SC \ge K_S^2 - (\delta_{\mathrm{alb}}(C)-1)$; then by Step 3 we have $K_SC \ge (\delta_{\mathrm{alb}}(C)-1)(4K_SC-1) \ge 4K_SC-1$ and thus $K_SC=0$. Then C is a -2 curve, $\delta_{\mathrm{alb}}(C)=q$ and $K_S^2 \le q-1$. The last equality contradicts the fact that $p_g(S) \ge q$ since S is of general type and $K_S^2 \ge 2p_g(S)$ since S is also irregular.

This concludes the proof of Theorem 1.1.

4. EXAMPLES

In this section we compute explicitly K_SC for some Albanese defective curves C contained in an irregular surface S. We examine products, symmetric covers and some double covers of abelian surfaces, namely most of the standard examples of irregular surfaces of general type. (One can of course also produce surfaces containing Albanese defective curves by taking suitable complete intersections in an abelian variety, but it is not clear how to compute explicitly K_SC and K_S^2 for such examples).

We use the notation introduced in the previous sections.

Example 4.1 (Curves in the product of two curves)**.** Let D be a curve of genus $g \geq 2$, let $S := D \times D$ and let $C \subset S$ be the diagonal. Here $A_C = J(D)$ and the quotient map $a_C \colon \operatorname{Alb}(S) = J(D) \times J(D) \to J(D)$ is defined by $(x, y) \mapsto x - y$, hence in this case the image of a_C is a surface. We have $\delta_{\mathrm{alb}}(C) = g$, $K_SC = 4(g-1) = K_S^2/2$. The map $S \to A_C$ maps S onto a surface. The situation is the same for graphs of automorphisms and similar for graphs of maps between curves.

If D_i, $i = 1, 2$, are curves of genus $g_i \geq 2$, $f \colon D_1 \to D_2$ is a non constant morphism and $C \subset S := D_1 \times D_2$ is the graph of f, then we also have $K_SC \leq K_S^2/2$, with equality holding iff f is étale.

Example 4.2 (Examples of curves with $\delta_{\mathrm{alb}}(C) = 1$ and $K_SC = K_S^2$)**.** Let E be an elliptic curve, set $A := E \times E$ and let $\Delta \subset A$ be the diagonal. Fix $x_1 \neq x_2 \in E$, set $B := \{x_1\} \times E + E \times \{x_1\} + \{x_2\} \times E + E \times \{x_2\}$, $P_{ij} = (x_i, x_j) \in A$, $i, j = 1, 2$. Choose $L \in \operatorname{Pic}(A)$ such that $2L \cong B$ and $L|_\Delta \not\equiv P_{11} + P_{22}$, and let $\pi \colon X \to A$ be the double cover associated to the relation $2L \equiv B$. The surface X has 4 singular points of type A_1, occurring over the points P_{ij}. The minimal resolution $S \to X$ is obtained by pulling back π to the blow up $\varepsilon \colon \hat{A} \to A$ of A at the points P_{ij} and normalizing. If we denote by $E_{ij} \subset \hat{A}$ the exceptional curve over P_{ij}, by B' the strict transform of B and by L' the line bundle $\varepsilon^* L(-\sum E_{ij})$, then $p \colon S \to \hat{A}$ is the double cover associated to the relation $2L' \equiv B'$. Hence the surface S is minimal of general type with $K_S^2 = 4$, $p_g(S) = q(S) = 2$. Let $\Delta' \subset \hat{A}$ be the strict transform of Δ and let $C = p^{-1}(\Delta')$. The double cover $C \to \Delta'$ is étale, given by the relation $2L'|_{\Delta'} \equiv 0$; since $L'|_{\Delta'}$ is non trivial, C is a smooth elliptic curve. Standard computations give $K_SC = 4 = K_S^2$.

Now pick $\eta \in \operatorname{Pic}^0(S)$ of finite order m such that $\eta|_C$ has also order m, let $S' \to S$ be the étale cover given by η and let $C' \subset S'$ be the preimage of C. Then C' is an elliptic curve with $C'K'_S = 4m = K_S^2$.

Example 4.3 (Curves in the symmetric square of a curve, I)**.** Let D be a curve of genus $q \geq 3$ and set $S := S^2D$. The surface S is minimal of general type with invariants $p_g(S) = q(q-1)/2$, $q(S) = q$, $K_S^2 = (q-1)(4q-9)$. Let σ be an involution of D; the curve $C := \{x = \sigma(x) | x \in D\} \subset S$ is a smooth curve isomorphic to D/σ. The pull back of C on $D \times D$ is the graph of σ, hence it has self intersection $2 - 2q$; it follows $C^2 = 1 - q$ and $K_SC = q - 1 + 2g - 2$, where g is the genus of C. By the Hurwitz formula we have $K_SC \leq 2q - 2$ with equality holding iff σ is base point free. The abelian variety A_C is isomorphic to the generalized Prym of the double cover $D \to D/\sigma$ and the map $a_C \colon S \to A_C$ is induced by the Abel-Prym map.

So, unless $q = 3$ and $g = 2$, $\delta_{\mathrm{alb}}(C) = g - q > 1$ and the image of a_C is a surface.

Example 4.4 (Curves in the symmetric square of a curve, II)**.** Let D, S and σ be as in Example 4.4 and assume in addition that $E := D/\sigma$ is an elliptic curve. Let $p\colon D \to E$ be the quotient map and let $f\colon S^2D \to E$ be defined by $(x, y) \mapsto x + y$. Let $c \in E$ and write $F_c := f^{-1}(c)$; the pull back $\tilde{F}_c$ of F_c to $D \times D$ is the preimage of $E_c := \{z + w = c\} \subset E \times E$ via the map $p \times p\colon D \times D \to E \times E$. Denote by B the branch locus of p and by $\mathrm{pr}_i\colon E \times E \to E$ the projection onto the i-th factor, $i = 1, 2$. The map $\tilde{F}_c \to E_c$ is a $\mathbb{Z}_2^2$-cover, which is the fiber product of two double covers branched on $\mathrm{pr}_i^* B$. By the Hurwitz formula B consists of $2q - 2$ distinct points and for $c \in E$ general the curve $\tilde{F}_c$ is smooth of genus $4q-3$. It is easy to check that the general $\tilde{F}_c$ meets the diagonal $\Delta_D \subset D\times D$ transversely at 8 points and therefore, again by the Hurwitz formula, it is smooth of genus $g := 2q - 1$.

Write $B = x_1 + \cdots + x_{2q-2}$. If, say, $x_1 = -x_2$ and $2x_1 \neq 0$, then $\tilde{F}_0$ has a node over the point $(x_1, -x_1)$ and a node over the point $(-x_1, x_1)$; these nodes are identified by the involution that exchanges the factors of $D \times D$, hence F_0 has a node. (If $x_1 = -x_1$, then $\tilde{F}_0$ has also a node over $(x_1, -x_1) = (x_1, x_1)$, but F_0 is smooth at the corresponding point). In a similar way one can arrange that F_0 has any number $r \leq q-1$ of nodes. For $r \leq q - 2$ the curve $\tilde{F}_0$ stays irreducible, hence F_0 is also irreducible. (For $r = q - 1$ it is possible to show that F_0 splits as the union of two smooth curves C_1, C_2 with $\delta_{\mathrm{alb}}(C_i) \geq 2$ for $i = 1, 2$ and $q \geq 4$.)

References

[Be1] A. Beauville, *Surfaces algébriques complexes*, Astérisque, no. 54, 1978.

[Be2] A. Beauville, *L'inégalité $p_g \geq 2q - 4$ pour les surfaces de type général*, appendix to O. Debarre, *Inégalités numériques pour les surfaces de type général*, Bull. Soc. Math. France **110** 3 (1982), 319–346.

[Ci] C. Ciliberto, *The bicanonical map for surfaces of general type*, Algebraic geometry - Santa Cruz 1995, 57–84, Proc. Sympos. Pure Math., **62**, Part 1, Amer. Math. Soc., Providence, RI, 1997.

[Lu] S.Y. Lu, *On surfaces of general type with maximal Albanese dimension*, J. Reine Angew. Math. **641** (2010), 163–175.

[LM] S.Y. Lu, Y. Miyaoka, *Bounding curves in algebraic surfaces by genus and Chern numbers*, Math. Res. Lett. **2**, no. 6 (1995), 663–676.

[Ma] M. Manetti, *Surfaces of Albanese general type and the Severi conjecture*, Math. Nach. **261–262** (2003), 105-122.

[Mi] Y. Miyaoka, *The orbibundle Miyaoka-Yau-Sakai inequality and an effective Bogomolov-McQuillan theorem*, Publ. Res. Inst. Math. Sci. **44** (2) (2008), 403–417.

[MP1] M. Mendes Lopes, R. Pardini *The geography of irregular surfaces*, in "Current developments in algebraic geometry", MSRI Publications, Volume 59 (2011).

[MP2] M. Mendes Lopes, R. Pardini, *Severi type inequalities for irregular surfaces with ample canonical class*, Comm. Math. Helv. 86, no. 2 (2011), 401–414.

[Pa] R. Pardini, *The Severi inequality $K^2 \geq 4\chi$ for surfaces of maximal Albanese dimension*, Invent. Math. **159** 3 (2005), 669 –672.

Margarida Mendes Lopes
Departamento de Matemática
Instituto Superior Técnico
Universidade Técnica de Lisboa
Av. Rovisco Pais
1049-001 Lisboa, PORTUGAL
mmlopes@math.ist.utl.pt

Rita Pardini
Dipartimento di Matematica
Università di Pisa
Largo B. Pontecorvo, 5
56127 Pisa, Italy
pardini@dm.unipi.it